

OSTEOPOROSIS ORDER FORM

PATIENT INFORMATION				
Last Name:	First Name:	MI DOB:		
HT: in WT: kg Birth Sex :() Male () Female Allergies: () NKDA, (Or):			
Provider Name	Contact Name	Contact Phone #		
NPI #:	Tax ID #:	Fax #:		
STATEMENT OF MEDICAL NECESSITY				
Primary Diagnosis: (ICD-10 CODE + DESCRIPTION)			
		Date of Diagnosis:		

PERTINENT MEDICAL HISTORY

Does patient have venous access?
YES NO If yes, what type MEDIPORT PIV PICC LINE OTHER:_

a) BIOSIMILAR EQUIVALENT SUBSTITUTION MAY APPLY

PRESCRIPTION ORDERS

SELECT	MEDICATION	DOSE	ROUTE	FREQUENCY	DURATION
	RECLAST (ZOLEDRONIC ACID)ADMINISTER OVER NO LESS THAN 15 MINUTESBUN, CREAT, AND CALCIUM LEVEL WITHIN 30 DAYS OF APPOINTMENTHOLD IF CALCIUM LEVELS < 8.5mg/dL	5 mg	IV	ONCE EVERY 12 MONTHS	1 Year
	PROLIA (DENOSUMAB) BUN, CREAT, CALCIUM LEVEL WITIN 90 DAYS OF THE APPOINTMENT HOLD IF CALCIUM LEVELS < <u>8.5mg/dL</u> or IONIZED CALCIUM LEVEL < <u>4.5mg/dL</u> or IF CRCL < <u>30 ML/MIN</u>	60 mg	Sub Q	ONCE EVERY 6 MONTHS	1 Year
	EVENITY BUN, CREAT, CALCIUM LEVEL WITIN 90 DAYS OF THE APPOINTMENT HOLD IF CALCIUM LEVELS < <u>8.5 mg/</u> dL or IONIZED CALCIUM LEVEL < <u>4.5 mg/dL</u> or IF CRCL < <u>30 ML/MIN</u>	210 mg	Sub Q	ONCE EVERY MONTH x 12	1 Year

LAB ORDERS: Calcium, BUN, Serum Creatinine will be drawn prior to administration is previous results not provided within 90 days of appointment.

SUPPORTING DOCUMENTATION FOR PATIENTS RECEIVING RECLAST, PROLIA, OR EVENITY:

- 1) OSTEOPOROSIS
- CALCIUM, BUN, AND SERUM CREATININE TO BE CHECKED WITHIN 30 DAYS FOR PATIENTS TAKING RECLAST OR 90 DAYS FOR PATIENTS TAKING PROLIA
- ORIGINAL BONE DENSITY/DEXA SCAN SUPPORTING THE DIAGNOSIS OF OSTEOPOROSIS
- H+P OR OFFICE NOTES LISTING THE DIAGNOSIS OF OSTEOPOROSIS IN THE PATIENT RECORD DATED WITHIN 1 YEAR PRIOR TO APPOINTMENT
- PRIOR/CURRENT MEDICATIONS USED TO TREAT THE DIAGNOSIS OF OSTEOPOROSIS MUST BE DOCUMENTED IN PATIENT'S MEDICAL RECORD (Examples: Oral calcium, Vitamin D, Bisphosphonates)
- 2) MEN AT HIGH RISK OF FRACTURE RECEIVING ANDROGEN DEPRIVATION THERAPY FOR NONMETASTATIC PROSTATE CANCER
- 3) TREATMENT TO INCREASE BONE MASS IN WOMEN AT HIGH RISK FOR FRACTURE RECEIVING AROMATASE INHIBITOR THERAPY FOR BREAST CANCER

*OSTEOPENIA IS NOT AN APPROVED DIAGNOSIS FOR PROLIA (DENOSUMAB). PATIENTS WITH IMPRESSIONS OF OSTEOPENIA MUST HAVE AN ORIGINAL BONE DENISTY RESULT OR DEXA SCAN SUPPORTING THE DIAGNOSIS OF OSTEOPOROSIS OR DOCUMENTATION OF A PREVIOUS FRAGILITY FRACTURE.

*PLEASE SUBMIT DOCUMENTATION OF ANY TRIED AND FAILED ORAL / INJECTIBLE MEDICATIONS ALONG WITH THE SUPPORTING DOCUMENTATION OF THE PATIENT RESPONSE / FAILURE TO TREATMENT.

*PROLIA, RECLAST, AND EVENITY ARE CONTRAINDICATED IN PATIENTS WITH HYPOCALCEMIA.

*EVENITY SHOULD NOT BE ADMINISTERED TO PATIENTS WHO HAVE A HISTORY OF STROKE OR MI (MYOCARDIAL INFARCTION) WITHIN THE LAST 12 MONTHS.

Physician's Signature _____ Date _____ Date _____

Signature must be clear and Legible	*Signature Must Be Clear and Legible	
-------------------------------------	--------------------------------------	--

Cosignature (If Required)_

*Signature Must Be Clear and Legible

Fax completed form, supporting documentation, facesheet, and insurance cards to the Outpatient Infusion Center at 1 (877) 249-1191.

Time_

Date_